

TransIT®-mRNA Transfection Kit

Quick Reference Protocol

Instructions for MIR 2225, 2250, 2251, 2255, 2256

Full protocol, SDS and Certificate of Analysis available at mirusbio.com/2250



SPECIFICATIONS

Storage	Store both <i>TransIT</i> ®-mRNA Reagent and mRNA Boost Reagent tightly capped at 4°C. Before each use , warm to room temperature and vortex gently.
Product Guarantee	1 year from the date of purchase, when properly stored and handled.

▶ RNA TRANSFECTION PROTOCOL



Full protocol and additional documentation available at mirusbio.com/2250

Fill in volumes below based on culture vessel used for transfection (Table 1).

A. Plate cells

1. Plate cells in ___ml complete growth medium (per well).

For adherent cells: Plate cells at a density of $0.8 - 3.0 \times 10^5$ cells/ml.

For suspension cells: Plate cells at a density of $2.5 - 5.0 \times 10^5$ cells/ml.

2. Culture overnight. Most cell types should be $\geq 80\%$ confluent on day of transfection.

B. Prepare *TransIT*®-mRNA Reagent:mRNA Boost:RNA complexes

1. Warm *TransIT*®-mRNA and mRNA Boost Reagents to room temperature and vortex gently.
2. Place ___ μ l of Opti-MEM® I Reduced-Serum Medium in a sterile tube.
3. Add ___ μ l RNA. Mix gently by pipetting.
4. Add ___ μ l of mRNA Boost Reagent. Mix gently by pipetting.
5. Add ___ μ l of *TransIT*®-mRNA Reagent. Mix gently by pipetting.
6. Incubate at room temperature for **2-5 minutes**.

C. Distribute complexes to cells

1. Add *TransIT*®-mRNA Reagent:mRNA Boost:RNA complex mixture drop-wise to different areas of the well.
2. Gently rock plate for even distribution of complexes.
3. Incubate 4-48 hours.
4. Harvest cells and assay as required.

Table 1. Recommended starting conditions

Culture vessel	24-well plate	12-well plate	6-well plate
Surface area	1.9 cm ²	3.8 cm ²	9.6 cm ²
Complete growth medium	0.5 ml	1 ml	2.5 ml
Serum-free medium	50 μ l	100 μ l	250 μ l
RNA (1 μ g/ μ l stock)	0.5 μ l	1 μ l	2.5 μ l
<i>TransIT</i> ®-mRNA Reagent	1 μ l	2 μ l	5 μ l
mRNA Boost Reagent	1 μ l	2 μ l	5 μ l

▶ Transfection Optimization

Determine the best *TransIT*®-mRNA:RNA and mRNA Boost:RNA ratio for each cell type. Start with 2 μ l of *TransIT*®-mRNA Reagent per 1 μ g of RNA. Vary the amount of *TransIT*®-mRNA Reagent from 1-3 μ l per 1 μ g RNA to find the optimal ratio. Vary the amount of mRNA Boost Reagent from 1-3 μ l per 1 μ g of RNA.

For additional optimization tips, see [full protocol](#).

Mirus Bio LLC

www.mirusbio.com | techsupport@mirusbio.com | U.S. Toll Free: 844.MIRUSBIO (844.647.8724) | Direct: +1.608.441.2852



Reagent Agent[®]

Reagent Agent[®] is an online tool designed to help determine the best solution for nucleic acid delivery based on in-house data, customer feedback and citations.

Learn more at: mirusbio.com/ra

©1996-2024 All rights reserved. Mirus Bio LLC. All trademarks are the property of their respective owners.

For terms and conditions, visit www.mirusbio.com

Rev0 100621